



## Advances in Dual Diagnosis

Trajectories of emotional symptoms in adolescence: impact on alcohol use

Louise Elizabeth Birrell, Nicola Clare Newton, Lexine Stapinski, Katrina Prior, Katrina Elizabeth Champion, Clare J. Mackie, Maree Teesson, Tim Slade,

### Article information:

To cite this document:

Louise Elizabeth Birrell, Nicola Clare Newton, Lexine Stapinski, Katrina Prior, Katrina Elizabeth Champion, Clare J. Mackie, Maree Teesson, Tim Slade, (2017) "Trajectories of emotional symptoms in adolescence: impact on alcohol use", *Advances in Dual Diagnosis*, Vol. 10 Issue: 4, pp.130-141, <https://doi.org/10.1108/ADD-08-2017-0014>

Permanent link to this document:

<https://doi.org/10.1108/ADD-08-2017-0014>

Downloaded on: 18 February 2018, At: 18:35 (PT)

References: this document contains references to 46 other documents.

To copy this document: [permissions@emeraldinsight.com](mailto:permissions@emeraldinsight.com)

The fulltext of this document has been downloaded 72 times since 2017\*

### Users who downloaded this article also downloaded:

(2017), "Psychosocial factors associated with adolescent substance use: a longitudinal investigation", *Advances in Dual Diagnosis*, Vol. 10 Iss 4 pp. 142-154 <<https://doi.org/10.1108/ADD-07-2017-0007>><https://doi.org/10.1108/ADD-07-2017-0007></a>

(2017), "Young construction workers: substance use, mental health, and workplace psychosocial factors", *Advances in Dual Diagnosis*, Vol. 10 Iss 4 pp. 155-168 <<https://doi.org/10.1108/ADD-08-2017-0013>><https://doi.org/10.1108/ADD-08-2017-0013></a>

Access to this document was granted through an Emerald subscription provided by emerald-srm:486125 []

### For Authors

If you would like to write for this, or any other Emerald publication, then please use our Emerald for Authors service information about how to choose which publication to write for and submission guidelines are available for all. Please visit [www.emeraldinsight.com/authors](http://www.emeraldinsight.com/authors) for more information.

### About Emerald [www.emeraldinsight.com](http://www.emeraldinsight.com)

Emerald is a global publisher linking research and practice to the benefit of society. The company manages a portfolio of more than 290 journals and over 2,350 books and book series volumes, as well as providing an extensive range of online products and additional customer resources and services.

Emerald is both COUNTER 4 and TRANSFER compliant. The organization is a partner of the Committee on Publication Ethics (COPE) and also works with Portico and the LOCKSS initiative for digital archive preservation.

\*Related content and download information correct at time of download.

# Trajectories of emotional symptoms in adolescence: impact on alcohol use

Louise Elizabeth Birrell, Nicola Clare Newton, Lexine Stapinski, Katrina Prior, Katrina Elizabeth Champion, Clare J. Mackie, Maree Teesson and Tim Slade

Louise Elizabeth Birrell, Nicola Clare Newton, Lexine Stapinski and Katrina Prior are all based at the Centre of Research Excellence in Mental Health and Substance Use, National Drug and Alcohol Research Centre, University of New South Wales, Sydney, Australia. Katrina Elizabeth Champion is a Postdoctoral Fellow at the National Drug and Alcohol Research Centre, University of New South Wales, Sydney, Australia. Clare J. Mackie is based at the Psychology & Neuroscience, Addictions Department, Institute of Psychiatry, Kings College London, London, UK. Maree Teesson and Tim Slade are both based at the Centre of Research Excellence in Mental Health and Substance Use, National Drug and Alcohol Research Centre, University of New South Wales, Sydney, Australia.

Received 23 August 2017  
Revised 29 September 2017  
2 October 2017  
Accepted 2 October 2017

This study is funded by the National Health and Medical Research Council (Grant No. APP1004744). The authors would like to acknowledge Associate Investigators on the grant as well as the schools, teachers, and students who have agreed to participate in this research. Louise Birrell was funded by an Australian Rotary Health PhD Scholarship.

## Abstract

**Purpose** – The purpose of this paper is to explore how different trajectories of emotional symptoms relate to alcohol use in adolescence.

**Design/methodology/approach** – In all, 431 participants (majority female), aged approximately 13 years at baseline were followed over three years and reported on their emotional symptoms and alcohol use. Latent class growth analyses explored different trajectories of emotional symptoms and regression models were run to relate these trajectories to alcohol use (full standard drink, and binge drinking) at 36-month follow-up (age 16 years).

**Findings** – While the majority of adolescents were best characterised by low-stable emotional symptoms, those with high-stable symptoms were more likely to have consumed a full standard drink of alcohol and binge drunk when aged 16 years.

**Research limitations/implications** – Findings highlight the importance of prevention and early intervention, particularly targeting adolescents with elevated stable emotional symptoms who were more likely to be using alcohol at 16 years of age.

**Originality/value** – The present study is one of the first longitudinal investigations into the use of alcohol by community adolescents with different emotional symptom trajectories.

**Keywords** Dual diagnosis, Trajectories, Development, Alcohol, Adolescence, Internalizing

**Paper type** Research paper

## Introduction

Problematic alcohol use and emotional disorders (anxiety and mood) cause significant disability and commonly co-occur (Kessler *et al.*, 2012; Teesson *et al.*, 2009). Compared to adult populations, relatively little research has been conducted exploring the development of this comorbidity in adolescents (Boschloo *et al.*, 2011; Hasin *et al.*, 2007). Furthermore, research examining emotional symptom and alcohol use development commonly fails to explore whether developmental trajectories of these behaviours are the same for all individuals. As such, longitudinal analysis examining both within, and between-person change during adolescence is needed to advance the field. The terms “emotional symptoms” and “emotional disorders” are used in this study as broadband terms capturing elements of both depression and anxiety symptomatology and disorders.

There is now substantial evidence that emotional symptom trajectories differ among individuals (Nandi *et al.*, 2009) and a more nuanced approach, which takes into account these differences rather than simply representing a mean growth rate for all individuals, is needed. This recognition has led to research attempting to identify subgroups of individuals that differ in their trajectories of emotional symptoms across adolescence (Dekker *et al.*, 2007; Rodriguez *et al.*, 2005; Stoolmiller *et al.*, 2005). Such studies typically utilise growth mixture modelling or latent class growth analysis (LCGA). Although these studies have used different age ranges and identify different numbers of emotional symptom trajectories, four typical trajectories are consistently observed in adolescence: low symptoms, high symptoms, moderate symptoms, and an

increasing trajectory (Brendgen *et al.*, 2005; Dekker *et al.*, 2007; Wiesner and Kim, 2006). When males are examined in isolation it is typical to observe a small number of adolescents characterised by decreasing symptoms (Stoolmiller *et al.*, 2005), while females are more likely to show increasing trajectories (Dekker *et al.*, 2007).

Research suggests that high emotional symptoms are also linked to increased alcohol use in adolescence (Measelle *et al.*, 2006; Sihvola *et al.*, 2008). For example, adolescents characterised by a high and persistent risk of experiencing an anxiety or depressive disorder have been found to be more likely to meet criteria for an alcohol use disorder (Olinio *et al.*, 2010). Yet to date, there has been no research examining how different emotional symptom trajectories relate to alcohol use in adolescence. In addition, previous studies examining alcohol use and emotional problems have tended to focus on one aspect of alcohol use, such as alcohol initiation or binge drinking, and failed to simultaneously explore the relationship between different drinking outcomes. Epidemiological evidence has shown that developmental pathways to alcohol use disorders are created well before problematic use of alcohol starts (Clark, 2004). Therefore, longitudinal research examining alcohol use at the lower end of the spectrum is needed to explore initial relationships between emotional symptoms and alcohol use when they first begin. Longitudinal evidence exploring how individuals with different trajectories of emotional symptoms consume alcohol over a critical adolescent period is needed to further understand the development of comorbid alcohol use and mental health problems and consider individual differences.

Building on existing research on emotional symptom trajectories, the aim of the current study is to conduct an investigation of how different trajectories of emotional symptoms relate to alcohol use (consuming a full standard drink of alcohol and binge drinking) over an important adolescent period (13-16 years). This is the first time this has been conducted in an Australian longitudinal sample. Based on previous literature, it is expected that the emotional symptom trajectories of adolescents will be characterised by four different groups: some starting at a high level and remaining high, some starting at a low level and remaining low, some experiencing increasing symptoms, and some showing middle-range stable symptoms. In addition, given the frequent comorbidity between emotional problems and alcohol use it is expected that adolescents with higher emotional symptoms will report higher alcohol use than those with lower symptom profiles.

## Method

### *Participants and procedure*

Data were drawn from the “Climate and Preventure” study, a cluster randomised controlled trial (RCT) evaluating the effectiveness of a prevention programme for alcohol and related harms in adolescents (for further details see Newton *et al.*, 2012). Only the control group of the RCT,  $n = 432$  from five independent (private) schools in Sydney, Australia, was used for the current study to avoid intervention effects confounding natural symptom trajectories. All Year 8 students attending participating schools in 2012 were invited to take part in the study. Only those students with parental consent (passive parental consent) and who gave active consent themselves were eligible to participate. Eligible participants completed a total of five online surveys over three years, including baseline, six-month, 12-month, 24-month, and 36-month follow-up post-baseline. The self-report questionnaire was completed by participants in a supervised (by project staff or teacher) classroom setting. Student data were linked over time using unique identifier codes to maintain confidentiality. One student was excluded from the analysis due to missing data on the emotional symptoms scale at each wave. The follow-up rate for the final sample ( $n = 431$ ) was 85 per cent at six-months, 91 per cent at 12-months, 86 per cent at 24-months, and 78 per cent at 36-month follow-up. Participants were 72 per cent female ( $n = 310$ ), mean age 13.38 years ( $SD = 0.44$ ) at baseline. All aspects of the study were approved by the Human Research Ethics Committee at the University of New South Wales.

### *Measures*

*Demographic information.* Participants were asked to report their age and gender.

*Assessment of emotional symptoms.* Emotional symptoms were measured through the five-item emotional symptoms subscale of the youth self-report Strengths and Difficulties Questionnaire (SDQ; Goodman, 2001). Participants rated items on a Likert scale ranging from 0 (not true) to 2 (certainly true) in relation to the previous six-months. Higher scores reflected greater emotional problems (range: 0-10). The emotional symptoms subscale is intended to capture internalising problems, such as anxiety and depression symptoms, and has good psychometric properties (Goodman, 2001). In the current study the internal reliability was acceptable to good across all study waves (all  $\alpha > 0.72$ ).

*Assessment of alcohol use.* Alcohol use at the 36-month follow-up was assessed via two drinking outcomes: whether students had consumed a full standard drink of alcohol (defined as containing ten grams of alcohol), and whether they had drunk alcohol to excess/"binged" (defined as consuming five or more standard drinks on one occasion), in the previous six months. A standard drink chart was provided to participants for reference. These measures were dichotomised as yes/no from continuous frequency measures, with the response option "never" coded "no", while any of the response options: "less than monthly", "monthly", "weekly", and "daily" were coded "yes". Although some information was lost through this process it was warranted given the overall low prevalence of alcohol use in the young community sample. Questions were modelled on those used in the School Health and Alcohol Harm Reduction Project "Patterns of Alcohol" index (McBride *et al.*, 2004), commonly used in adolescent populations (Conrod *et al.*, 2008; Newton *et al.*, 2009; Vogl *et al.*, 2009).

*Externalising symptoms.* Externalising symptoms at the 36-month follow-up were assessed through the ten-item externalising subscale of the SDQ. Participants were asked to respond to items on a three-point Likert scale ranging from 0 (not true) to 2 (certainly true) in relation to the previous six months. The internal reliability of the externalising subscale was good in the current study ( $\alpha = 0.79$ ).

### **Statistical analysis**

The analysis strategy consisted of two parts:

1. LCGA modelled the different trajectories of emotional symptoms in adolescence.
2. Regression models were run to compare alcohol use at 36-month follow-up, with emotional symptom trajectory classes as predictors. Based on the study hypotheses, those with high emotional symptoms were compared to other adolescents.

### **Trajectories of emotional symptoms**

A number of unconditional latent class growth models were run in Mplus 7.4 (Muthén and Muthén, 1998-2010). Missing data at each follow-up wave were accommodated through full information maximum-likelihood estimation, which estimates parameters using all available data. As descriptive plots indicated some positive skew in emotional symptom data, models were estimated using maximum likelihood with robust standard errors (Little and Rubin, 1987) which is robust to violations of multivariate normality (Schafer and Graham, 2002). Models were run with multiple starts to ensure the robustness of the log likelihood and avoid a false maximum likelihood optima (Hipp and Bauer, 2006; Morin *et al.*, 2011). One to seven class solutions were computed. Recommendations of Ram and Grimm (2009) were followed to select the best fitting model, which included an inspection of the Bayesian Information Criteria (BIC), Akaike Information Criteria (AIC), Vuong-Lo-Mendell-Rubin Likelihood Ratio Test and Adjusted Lo-Mendell-Rubin likelihood. After identifying the best fitting model, individuals were assigned to their most likely class based on their posterior probability values (average latent class probabilities were all  $> 0.83$ ).

### **Examining alcohol use by emotional trajectories**

Multiple logistic regression models were run in SPSS 23. All regression models were run with probability weightings applied, taking into account inherent uncertainty regarding class

membership in latent class modelling. Separate models were run for the following outcomes: consumed a full standard drink of alcohol, and binge drank. Unconditional (unadjusted) models were run comparing those characterised by high-stable emotional symptoms to the other classes. A second set of models (adjusted) was run with the addition of the following covariates added in one block: gender, baseline alcohol use, and externalising symptoms. Previous literature has established a link between externalising symptoms (e.g. aggressive, oppositional, hyperactive symptoms) and alcohol use in adolescence (Jun *et al.*, 2015; King *et al.*, 2004). It has also consistently been found that females experience more emotional symptoms in adolescence compared to males (Garber *et al.*, 2002; Ge *et al.*, 2006) and that males use more alcohol (Nolen-Hoeksema, 2004). Therefore, these potential confounders were included in the adjusted models.

## Results

### Latent class growth model selection

A series of unconditional latent class growth models ranging from one to seven classes were estimated with: an intercept and linear slope (fit indices displayed in Table A1), and an intercept, linear slope, and quadratic slope (fit indices shown in Table I).

### Model fit

Model fit indices showed that on balance the best fitting and most parsimonious model was the four-class model with an intercept, linear and quadratic growth term (AIC = 7602.02, BIC = 7683.39, Entropy = 0.79, LMR-ALRT = 50.01,  $p < 0.05$ , BLRT  $p$ -value  $< 0.01$ ). Quadratic growth parameters were observed in a number of models; therefore, models including a linear and quadratic growth term were examined. The four-class model showed satisfactory entropy with a value of 0.79, indicating adequate-good separation between classes.

The fit indices for the four-class quadratic growth model are presented in Table I. On balance, the four-class model was selected as providing the best fit to the data as well being the most parsimonious.


### Trajectory classes

Figure 1 displays emotional symptom trajectories by latent class. There were 195 (45 per cent) adolescents best characterised by low initial levels of emotional symptoms (intercept = 1.21,  $p < 0.01$ ) that remained stable and low over the three years (slope = 0.21,  $p = 0.34$ ), named the low-stable class. In all, 17 (4 per cent) adolescents were best characterised by a quadratic growth trajectory of initial rapid increase in emotional symptoms that then decreased between age 15 and 16 years, named the rapid-increase class (intercept = 0.41,  $p = 0.46$ ; slope = 6.57,

**Table I** Fit indices for latent class growth analysis models with linear and quadratic growth term

	Parameters	AIC	BIC	Entropy	LMR-ALRT	BLRT $p$ -value
1 Class	8	8,235.05	8,267.60	–	–	–
2 Classes	12	7,773.29	7,822.11	0.78	451.17**	$< 0.01$
3 Classes	16	7,646.09	7,711.18	0.74	129.85**	$< 0.01$
4 Classes	20	7,602.02	7,683.39	0.79	50.01*	$< 0.01$
5 Classes	24	7,585.27	7,682.92	0.75	23.77	$< 0.01$
6 Classes	28	7,577.28	7,691.20	0.77	15.36	$< 0.01$
7 Classes	32	7,568.61	7,698.80	0.77	16.02	$< 0.01$

**Notes:** AIC, Akaike Information Criterion; BIC, Bayesian Information Criterion; LMR-LRT, Lo-Mendall-Rubin adjusted Likelihood Ratio Test; BLRT, Bootstrapped Likelihood Ratio Test. Classes beyond the seven class model were not explored as there were less than ten people in an individual class in the seven class model, which is the suggested minimum for interpretability. \* $p < 0.05$ ; \*\* $p < 0.001$

**Figure 1** Four-class latent class growth analysis of trajectories of emotional symptoms

$p < 0.01$ ; quadratic =  $-1.71$ ,  $p < 0.01$ ). In total, 57 (13 per cent) adolescents had high initial levels of emotional symptoms that remained stable (intercept =  $5.99$ ,  $p < 0.01$ ; slope =  $0.65$ ,  $p = 0.15$ ), named the high-stable class. Finally, 162 (38 per cent) adolescents were in a class named the middle-range class, with initial middle-range emotional symptoms that remained stable (intercept =  $3.57$ ,  $p < 0.01$ ; slope =  $-0.09$ ,  $p = 0.80$ ).

#### Alcohol use by latent class

Table II reports alcohol use at baseline and 36-month follow-up by latent trajectory class.

#### Full standard drink of alcohol

Logistic regression results displayed in Table III indicated significant differences by trajectory class in relation to a full standard drink of alcohol at 36-month follow-up. Specifically, adolescents with high-stable emotional symptoms were 3.3 times (95 per cent CI [1.5, 7.4]) more likely to report having consumed a full standard drink of alcohol, compared to adolescents in the middle-range

**Table II** Percentage of participants who had used alcohol in the past six months at baseline and 36-month follow-up by emotional symptoms latent class membership

	Trajectory class			
	Class 1 – low-stable (%)	Class 2 – rapid-increase (%)	Class 3 – high-stable (%)	Class 4 – middle-range (%)
<i>Baseline</i>				
Had a full drink	6	6	12	7
Binged	3	6	5	4
<i>36-month follow-up</i>				
Had a full drink	59	36	77	48
Binged	40	29	51	32

**Table III** Multiple logistic regression models predicting full serve of alcohol in past six months at 36-month follow-up, with latent class group comparisons

Variable	Model 1 <sup>a</sup>					Model 2 <sup>b</sup>				
	$\beta$	SE	$p$	OR	95% CI	$\beta$	SE	$p$	OR	95% CI
Trajectory class comparisons										
High-stable vs middle-range	1.20	0.41	<0.01	3.32	1.49-7.40	1.32	0.44	<0.01	3.73	1.57-8.85
High-stable vs rapid-increase	1.47	0.69	0.03	4.37	1.14-16.80	1.84	0.74	0.01	6.32	1.50-26.67
High-stable vs low-stable	0.74	0.40	0.06	2.09	0.96-4.55	0.68	0.44	0.12	1.98	0.84-4.68
Gender						0.39	0.29	0.18	1.47	0.84-2.60
Baseline ever had full drink						1.55	0.47	<0.01	4.74	1.88-11.94
Externalising symptoms						<0.01	0.04	0.81	1.01	0.94-1.09

Notes: <sup>a</sup>Model 1 was unadjusted and looked at the effect of latent class on alcohol use only,  $n = 324$ ; <sup>b</sup>Model 2 adjusted for the covariates of gender, ever had full drink at baseline and externalising scores on the SDQ (at 36-month f/u),  $n = 309$ .  $\beta$  – regression coefficient; Gender – males as reference category

class ( $p < 0.01$ ). This remained significant when controlling for covariates (OR = 3.7, 95 per cent CI [1.6, 8.9],  $p < 0.01$ ). Adolescents with high-stable emotional symptoms were also 4.4 times more likely to have consumed a full standard drink of alcohol compared to those with a rapidly increasing trajectory (OR = 4.4, 95 per cent CI [1.1, 16.8],  $p = 0.03$ ). This remained significant when controlling for all covariates (OR = 6.3, 95 per cent CI [1.5, 26.7],  $p = 0.01$ ). There were no significant differences when comparing those in the high-stable trajectory to those in the low-stable emotional trajectory. Baseline alcohol use was the only significant covariate (OR = 4.7, 95 per cent CI [1.9, 11.9],  $p < 0.01$ ).

### Binge drinking

Table IV displays results in relation to binge drinking. Adolescents with a high-stable emotional symptom trajectory were 2.3 times more likely to report binge drinking at 36-month follow-up, compared to adolescents with middle-range emotional symptoms (OR = 2.3, 95 per cent CI [1.1, 4.7],  $p = 0.03$ ). This finding remained significant when controlling for covariates (OR = 2.2, 95 per cent CI [1.01, 4.8],  $p < 0.05$ ). There were no statistically significant differences comparing those in the high class to others in either the unadjusted or adjusted models. Externalising symptoms were the only significant covariate (OR = 1.1, 95 per cent CI [1.0-1.2],  $p = 0.01$ ).

### Discussion

This study is one of the first to investigate how alcohol use relates to trajectories of emotional symptoms in adolescence. Longitudinal data and sophisticated analytic methods were utilised to explore individual differences in change over time in a community sample of adolescents.

**Table IV** Multiple logistic regression models predicting binge drinking (five or more drinks) in past six months at 36-month follow-up, with latent class group comparisons

Variable	Model 1 <sup>a</sup>					Model 2 <sup>b</sup>				
	$\beta$	SE	$p$	OR	95% CI	$\beta$	SE	$p$	OR	95% CI
Trajectory class comparisons										
High-stable vs middle-range	0.81	0.38	0.03	2.25	1.0-4.71	0.79	0.40	<0.05	2.22	1.01-4.82
High-stable vs rapid-increase	0.72	0.69	0.30	2.05	0.53-7.87	0.77	0.72	0.28	2.16	0.53-8.80
High-stable vs low-stable	0.41	0.36	0.25	1.50	0.75-3.01	0.26	0.39	0.51	1.29	0.60-2.75
Gender						-0.06	0.29	0.89	0.94	0.53-1.66
Baseline ever binged						0.98	0.80	0.22	2.68	0.56-12.70
Externalising symptoms						0.09	0.04	0.02	1.10	1.02-1.18

Notes: <sup>a</sup>Model 1 was unadjusted and looked at the effect of latent class on alcohol use only,  $n = 324$ ; <sup>b</sup>Model 2 adjusted for the covariates of gender, ever binge drunk at baseline and externalising scores on the SDQ (at 36-month f/u),  $n = 309$ .  $\beta$  – regression coefficient; Gender – males as reference category

As expected, there was heterogeneity in the trajectories of emotional symptoms between 13 and 16 years, best characterised by four trajectories: low-stable, middle-range, high-stable, and rapidly increasing. In line with previous research, the majority of adolescents experienced low-stable emotional symptoms (45 per cent) or stable middle-range symptoms (38 per cent) over the three years. A substantial portion (13 per cent) started with relatively high emotional symptoms that remained high, while a minority (4 per cent) showed a rapidly increasing trajectory of emotional symptoms, demonstrating a non-linear trend over time.

These four observed trajectories are in line with previous research that has applied growth modelling to depression symptoms in adolescence (Brendgen *et al.*, 2005; Wiesner and Kim, 2006). The rapid-increasing trajectory observed in the current sample is also consistent with increasing trajectories observed in a previous adolescent female sample (Dekker *et al.*, 2007). However, one study observed a decreasing trajectory among males (Stoolmiller *et al.*, 2005), which was not observed in the current sample. The decreasing trajectory observed by Stoolmiller *et al.* (2005) was seen in relation to an older sample (aged 15-24 years) of males. The lack of a decreasing trajectory in the current sample may be reflective of the younger predominately female sample in the current study. It should be noted there was a relatively small number of individuals characterised by the rapidly increasing trajectory class ( $n = 17$ ). The estimates calculated and wide confidence intervals for this class in the regression models point to imprecise prediction of alcohol use for individuals characterised by this trajectory. Therefore, replication of this trajectory and its relationship to alcohol use with a larger sample size is needed.

In relation to alcohol use at 16 years of age, those with high-stable emotional symptoms were more likely to have consumed a full standard drink of alcohol in comparison to adolescents with middle-range emotional symptoms and those with a rapidly increasing trajectory. They were also more likely to have engaged in binge drinking when compared to those in the middle-range, but no different to those in low-stable or increasing-trajectory classes. These differences remained significant over and above the effect of gender, baseline drinking, and externalising symptoms. These findings are consistent with the literature linking heightened emotional symptoms and disorders to an increased chance of using alcohol and experiencing alcohol use problems (Reinke *et al.*, 2012) and extends these findings to adolescence. Several theoretical models can be used to explain the comorbidity between emotional symptoms and alcohol use. The “self-medication hypothesis”, Khantzian (1997) for example, proposes that there is a direct causal pathway between emotional symptoms and alcohol use. That is, adolescents with emotional problems drink alcohol to treat distressing emotional symptoms and problematic alcohol use develops over time. Similarly, “the tension reduction hypothesis” posits that people use alcohol to help to relieve symptoms of anxiety (Greeley and Oei, 1999). However, research among adolescents has not been able to directly determine whether young people use alcohol to self-medicate or reduce tension. Another possible explanation is that a range of common factors, including biological, environmental, and psychological factors, explain the comorbidity between emotional symptoms and alcohol use, in that some individuals are more likely to develop both.

The current study has significant implications for alcohol and mental health prevention. The finding that those with the highest emotional symptoms were the most likely to be involved with alcohol by age 16 years suggests that preventative efforts should target both emotional symptoms and alcohol use during early adolescence. One example of such an approach is the online school-based “Climate Schools Combined” programme, an integrated intervention to prevent both substance use and mental health problems among secondary school students, which is currently under evaluation (Teesson *et al.*, 2014). This universal approach (i.e. delivered to all students regardless of risk) aims to equip students with knowledge and skills in early adolescence, before young people typically initiate alcohol use and prior to the onset of mental health symptoms. Importantly, this online intervention can be readily implemented by teachers, without prior training. Alternatively, selective interventions that target “high-risk” adolescents, such as youth already exhibiting high emotional symptoms in early adolescence, may also be beneficial in reducing the shared risk factors for comorbid

alcohol use and mental health problems in later adolescence and adulthood. Possible delivery modes for a selective intervention include school or community settings; however, teacher or facilitator training would first be necessary. The fact that the next most likely drinkers were adolescents with low-stable emotional symptoms indicates that for some adolescents, factors other than emotional symptoms drive their drinking. It has been shown that adolescents with different motives for using alcohol, e.g. drinking to cope or drinking for social enhancement, show different drinking patterns (Stapinski *et al.*, 2016). Future research could extend this work by looking at different drinking motives for adolescents with different levels of emotional symptoms and develop tailored intervention approaches accordingly. Taken together, the present findings suggest that future prevention research may benefit from addressing an array of factors, including drinking motives and emotional symptoms, and should target both “low-risk” and “high-risk” students, such as those with high-stable emotional symptoms in early adolescence.

These findings reinforce the importance of early and effective prevention of alcohol use and mental health symptoms, as unrecognised emerging problems have the potential to lead to significant problems in late adolescence and early adulthood, including anxiety, mood and alcohol use disorders. These disorders are associated with significant burden of disease and pose huge social and economic challenges to society, through substantial health care costs and loss of participation in the workforce (Bloom *et al.*, 2011; Gore *et al.*, 2011; Hu, 2006; Whiteford *et al.*, 2013). Additionally, early onset alcohol use and mental disorders are associated with a more difficult life course, including unemployment (Paljarvi *et al.*, 2015) and lower adult income (Wilson *et al.*, 2015).

### Study strengths

Key strengths of this study include the examination of within-individual and non-linear patterns of change (non-uniform change e.g. rapid during some periods and slower in others), the developmental approach taken, and longitudinal study design. The use of a general community sample means results more accurately reflect the natural developmental progression of adolescent mental health problems compared to if a clinical sample had been used. Finally, this study examined alcohol use via two separate drinking outcomes, using specific measurements that were explained to participants in standard units where appropriate. This led to a higher level of precision than has been used in previous studies.

### Limitations

Conclusions drawn from the present study should be viewed in light of the following limitations. First, the fact that the present sample was predominately made up of females (72 per cent) attending independent schools in Sydney, may limit generalisability. Sample size restrictions also meant analyses could not be conducted separately for males and females and future research with larger samples are needed to confirm that the findings are equally applicable to males and females. While beyond the scope of this paper, future research could also examine gender as a moderator of the relationship between emotional symptoms and different levels of alcohol use. Adolescents were also asked to self-report emotional symptoms and alcohol use, which can be influenced by self-report biases, such as social desirability. However, good reliability and validity of self-reported substance use has been demonstrated (Del Boca and Darkes, 2003; Ramo *et al.*, 2012), especially when confidentiality is assured and when students self-administer surveys online (Brener *et al.*, 2003), both of which occurred in this study.

Finally, it has been argued that the application of trajectory modelling to characterise individual differences has consistently produced four prototypical trajectory classes, despite varying age ranges of samples and observation periods: a consistently low group, an increasing group, a decreasing group, and a consistently high group (Sher *et al.*, 2011). While this pattern has been found in relation alcohol use trajectories, there have been numerous studies of psychopathology trajectories that did not find this pattern. However, three of the typical

four trajectories were observed in the current study, suggesting that the generalisation of these results should be carried out with caution and the further validation of these four classes is required in other samples to confirm the validity of the class solution found in the present study.

## Conclusions

The present study is the first longitudinal investigation into the use of alcohol by community adolescents with different emotional symptom trajectories. The different trajectories of emotional symptoms are in line with previous literature and add further confirmation that emotional symptoms are heterogeneous. The novel aspect of this study was the discovery that these trajectories are related to different rates of alcohol use by age 16 years. This suggests interventions targeting both emotional problems and alcohol use may see more benefits than targeting either alone, especially among individuals with higher emotional symptoms. In addition, the relatively high proportion of adolescents with low-stable emotional symptoms using alcohol also suggests that other risk factors need to be included into prevention programmes to reduce these problems in the population.

## References

- Bloom, D.E., Cafiero, E.T., Jané-Llopis, E., Abrahams-Gessel, S., Bloom, L.R., Fathima, S., Feigl, A.B., Gaziano, T., Mowafi, M., Pandya, A., Prettnner, K., Rosenberg, L., Seligman, B., Stein, A.Z. and Weinstein, C. (2011), "The global economic burden of noncommunicable diseases".
- Boschloo, L., Vogelzangs, N., Smit, J.H., van den Brink, W., Veltman, D.J., Beekman, A.T. and Penninx, B.W. (2011), "Comorbidity and risk indicators for alcohol use disorders among persons with anxiety and/or depressive disorders: findings from the Netherlands Study of Depression And Anxiety (NESDA)", *Journal of Affective Disorders*, Vol. 131 Nos 1/3, pp. 233-42.
- Brendgen, M., Wanner, B., Morin, A.J. and Vitaro, F. (2005), "Relations with parents and with peers, temperament, and trajectories of depressed mood during early adolescence", *Journal of Abnormal Child Psychology*, Vol. 33 No. 5, pp. 579-94.
- Brener, N.D., Billy, J.O. and Grady, W.R. (2003), "Assessment of factors affecting the validity of self-reported health-risk behavior among adolescents: evidence from the scientific literature", *Journal of Adolescent Health*, Vol. 33 No. 6, pp. 436-57.
- Greeley, J. and Oei, T. (1999), "Alcohol and tension reduction: an update on research and theory", in Blane, H.T. and Leonard, K.E. (Eds), *Psychological Theories of Drinking and Alcoholism*, Guilford Press, New York, NY, pp. 14-54.
- Clark, D.B. (2004), "The natural history of adolescent alcohol use disorders", *Addiction*, Vol. 99 No. 1, pp. 5-22.
- Conrod, P., Castellanos, N. and Mackie, C. (2008), "Personality-targeted intervention delay the growth of adolescent drinking and binge drinking", *Journal of Child Psychology and Psychiatry*, Vol. 49 No. 2, pp. 181-90.
- Dekker, M.C., Ferdinand, R.F., van Lang, N.D., Bongers, I.L., van der Ende, J. and Verhulst, F.C. (2007), "Developmental trajectories of depressive symptoms from early childhood to late adolescence: gender differences and adult outcome", *Journal of Child Psychology and Psychiatry*, Vol. 48 No. 7, pp. 657-66.
- Del Boca, F.K. and Darkes, J. (2003), "The validity of self-reports of alcohol consumption: state of the science and challenges for research", *Addiction*, Vol. 98 No. S2, pp. 1-12.
- Garber, J., Keiley, M.K. and Martin, C. (2002), "Developmental trajectories of adolescents' depressive symptoms: predictors of change", *Journal of Consulting and Clinical Psychology*, Vol. 70 No. 1, pp. 79-95.
- Ge, X., Natsuaki, M.N. and Conger, R.D. (2006), "Trajectories of depressive symptoms and stressful life events among male and female adolescents in divorced and nondivorced families", *Development and Psychopathology*, Vol. 18 No. 1, pp. 253-73.

- Goodman, R. (2001), "Psychometric properties of the Strengths and Difficulties Questionnaire", *Journal of the American Academy of Child & Adolescent Psychiatry*, Vol. 40 No. 11, pp. 1337-45.
- Gore, F.M., Bloem, P.J., Patton, G.C., Ferguson, J., Joseph, V., Coffey, C., Sawyer, S.M. and Mathers, C.D. (2011), "Global burden of disease in young people aged 10-24 years: a systematic analysis", *Lancet*, Vol. 377 No. 9783, pp. 2093-102.
- Hasin, D.S., Stinson, F.S., Ogburn, E. and Grant, B.F. (2007), "Prevalence, correlates, disability, and comorbidity of DSM-IV alcohol abuse and dependence in the United States: results from the national epidemiologic survey on alcohol and related conditions", *Archives of General Psychiatry*, Vol. 64 No. 7, pp. 830-42.
- Hipp, J.R. and Bauer, D.J. (2006), "Local solutions in the estimation of growth mixture models", *Psychological Methods*, Vol. 11 No. 1, pp. 36-53.
- Hu, T. (2006), "Perspectives: an international review of the national cost estimates of mental illness, 1990-2003", *The Journal of Mental Health Policy and Economics*, Vol. 9 No. 1, pp. 3-13.
- Jun, H.J., Sacco, P., Bright, C.L. and Camlin, E.A. (2015), "Relations among internalizing and externalizing symptoms and drinking frequency during adolescence", *Substance Use and Misuse*, Vol. 50 No. 14, pp. 1814-25.
- Kessler, R.C., Avenevoli, S., McLaughlin, K.A., Green, J.G., Lakoma, M.D., Petukhova, M., Pine, D.S., Sampson, N.A., Zaslavsky, A.M. and Merikangas, K.R. (2012), "Lifetime co-morbidity of DSM-IV disorders in the US National comorbidity survey replication adolescent supplement (NCS-A)", *Psychological Medicine*, Vol. 42 No. 9, pp. 1997-2010.
- Khantzian, E.J. (1997), "The self-medication hypothesis of substance use disorders: a reconsideration and recent applications", *Harvard Review of Psychiatry*, Vol. 4 No. 5, pp. 231-44.
- King, S.M., Iacono, W.G. and McGue, M. (2004), "Childhood externalizing and internalizing psychopathology in the prediction of early substance use", *Addiction*, Vol. 99 No. 12, pp. 1548-59.
- Little, R.J.A. and Rubin, D.B. (1987), *Statistical Analysis with Missing Data*, Wiley, Hoboken, NJ.
- McBride, N., Farrington, F., Midford, R., Meuleners, L. and Phillips, M. (2004), "Harm minimisation in school drug education: final results of the school health and alcohol harm reduction project (SHAHRP)", *Addiction*, Vol. 99 No. 3, pp. 278-91.
- Measelle, J.R., Stice, E. and Hogansen, J.M. (2006), "Developmental trajectories of co-occurring depressive, eating, antisocial, and substance abuse problems in female adolescents", *Journal of Abnormal Psychology*, Vol. 115 No. 3, pp. 524-38.
- Morin, A.J.S., Maiano, C., Nagengast, B., Marsh, H.W., Morizot, J. and Janosz, M. (2011), "General growth mixture analysis of adolescents' developmental trajectories of anxiety: the impact of untested invariance assumptions on substantive interpretations", *Structural Equation Modeling – A Multidisciplinary Journal*, Vol. 18 No. 4, pp. 613-48.
- Muthén, L.K. and Muthén, B.O. (1998-2010), *Mplus User's Guide*, 6th ed., Muthén & Muthén, Los Angeles, CA.
- Nandi, A., Beard, J.R. and Galea, S. (2009), "Epidemiologic heterogeneity of common mood and anxiety disorders over the lifecourse in the general population: a systematic review", *BMC Psychiatry*, Vol. 9 No. 31.
- Newton, N.C., Vogl, L.E., Teesson, M. and Andrews, G. (2009), "CLIMATE schools: alcohol module: cross-validation of a school-based prevention programme for alcohol misuse", *Australian & New Zealand Journal of Psychiatry*, Vol. 43 No. 3, pp. 201-7.
- Newton, N.C., Teesson, M., Barrett, E.L., Slade, T. and Conrod, P.J. (2012), "The CAP study, evaluation of integrated universal and selective prevention strategies for youth alcohol misuse: study protocol of a cluster randomized controlled trial", *BMC Psychiatry*, Vol. 12 No. 118.
- Nolen-Hoeksema, S. (2004), "Gender differences in risk factors and consequences for alcohol use and problems", *Clinical Psychology Review*, Vol. 24 No. 8, pp. 981-1010.
- Olino, T.M., Klein, D.N., Lewinsohn, P.M., Rohde, P. and Seeley, J.R. (2010), "Latent trajectory classes of depressive and anxiety disorders from adolescence to adulthood: descriptions of classes and associations with risk factors", *Comprehensive Psychiatry*, Vol. 51 No. 3, pp. 224-35.

- Paljarvi, T., Martikainen, P., Pensola, T., Leinonen, T., Herttua, K. and Makela, P. (2015), "Life course trajectories of labour market participation among young adults who experienced severe alcohol-related health outcomes: a retrospective cohort study", *PLoS One*, Vol. 10 No. 5.
- Ram, N. and Grimm, K.J. (2009), "Growth mixture modeling: a method for identifying differences in longitudinal change among unobserved groups", *International Journal of Behavioral Development*, Vol. 33 No. 6, pp. 565-76.
- Ramo, D.E., Liu, H. and Prochaska, J.J. (2012), "Reliability and validity of young adults' anonymous online reports of marijuana use and thoughts about use", *Psychology of Addictive Behaviors*, Vol. 26 No. 4, pp. 801-11.
- Reinke, W.M., Eddy, J.M., Dishion, T.J. and Reid, J.B. (2012), "Joint trajectories of symptoms of disruptive behavior problems and depressive symptoms during early adolescence and adjustment problems during emerging adulthood", *Journal of Abnormal Child Psychology*, Vol. 40 No. 7, pp. 1123-36.
- Rodriguez, D., Moss, H.B. and Audrain-McGovern, J. (2005), "Developmental heterogeneity in adolescent depressive symptoms: associations with smoking behavior", *Psychosomatic Medicine*, Vol. 67 No. 2, pp. 200-10.
- Schafer, J.L. and Graham, J.W. (2002), "Missing data: our view of the state of the art", *Psychological Methods*, Vol. 7 No. 2, pp. 147-77.
- Sher, K.J., Jackson, K.M. and Steinley, D. (2011), "Alcohol use trajectories and the ubiquitous cat's cradle: cause for concern?", *Journal of Abnormal Psychology*, Vol. 120 No. 2, pp. 322-35.
- Sihvola, E., Rose, R.J., Dick, D.M., Pulkkinen, L., Marttunen, M. and Kaprio, J. (2008), "Early-onset depressive disorders predict the use of addictive substances in adolescence: a prospective study of adolescent Finnish twins", *Addiction*, Vol. 103 No. 12, pp. 2045-53.
- Stapinski, L.A., Edwards, A.C., Hickman, M., Araya, R., Teesson, M., Newton, N.C., Kendler, K.S. and Heron, J. (2016), "Drinking to cope: a latent class analysis of coping motives for alcohol use in a large cohort of adolescents", *Prevention Science*, Vol. 17 No. 5, pp. 584-94.
- Stoolmiller, M., Kim, H.K. and Capaldi, D.M. (2005), "The course of depressive symptoms in men from early adolescence to young adulthood: identifying latent trajectories and early predictors", *Journal of Abnormal Psychology*, Vol. 114 No. 3, pp. 331-45.
- Teesson, M., Slade, T. and Mills, K. (2009), "Comorbidity in Australia: findings of the 2007 national survey of mental health and wellbeing", *Australian & New Zealand Journal of Psychiatry*, Vol. 43 No. 7, pp. 606-14.
- Teesson, M., Newton, N.C., Slade, T., Chapman, C., Allsop, S. and Hides, L. (2014), "The CLIMATE schools combined study: a cluster randomised controlled trial of a universal Internet-based prevention program for youth substance misuse, depression and anxiety", *BMC Psychiatry*, Vol. 14 No. 32.
- Vogl, L., Teesson, M., Andrews, G., Bird, K., Steadman, B. and Dillon, P. (2009), "A computerised harm minimisation prevention program for alcohol misuse and related harms: randomised controlled trial", *Addiction*, Vol. 104 No. 4, pp. 564-75.
- Whiteford, H., Degenhardt, L., Rehm, J., Baxter, A.J., Ferrari, A.J., Erskine, H.E., Charlson, F.J., Norman, R.E., Flaxman, A.D., Johns, N., Burstein, R., Murray, C.J. and Vos, T. (2013), "Global burden of disease attributable to mental and substance use disorders: findings from the global burden of disease study 2010", *The Lancet*, Vol. 382 No. 9904, pp. 1575-86.
- Wiesner, M. and Kim, H.K. (2006), "Co-occurring delinquency and depressive symptoms of adolescent boys and girls: a dual trajectory modeling approach", *Developmental Psychology*, Vol. 42 No. 6, pp. 1220-35.
- Wilson, S., Hicks, B.M., Foster, K.T., McGue, M. and Iacono, W.G. (2015), "Age of onset and course of major depressive disorder: associations with psychosocial functioning outcomes in adulthood", *Psychological Medicine*, Vol. 45 No. 3, pp. 505-14.

## Appendix

**Table A1** Fit indices for latent class growth analysis models with linear growth term

	<i>Parameters</i>	<i>AIC</i>	<i>BIC</i>	<i>Entropy</i>	<i>LMR-ALRT</i>	<i>BLRT p-value</i>
1 Class	7	8,233	8,262	–	–	–
2 Classes	10	7,774	7,815	0.78	441.37**	< 0.01
3 Classes	13	7,646	7,699	0.75	127.16**	< 0.01
4 Classes	16	7,633	7,699	0.77	17.53	0.71
5 Classes	19	7,621	7,638	0.69	17.02	< 0.01
6 Classes	22	7,620	7,709	0.69	7.50	0.16
7 Classes <sup>a</sup>	25	7,616	7,718	0.67	8.88	0.11

**Notes:** AIC, Akaike Information Criterion; BIC, Bayesian Information Criterion; LMR-LRT, Lo-Mendall-Rubin adjusted Likelihood Ratio Test; BLRT, Bootstrapped Likelihood Ratio Test. <sup>a</sup>Classes beyond the seven class model were not explored as there were less than ten people in an individual class in the seven class model, which is the suggested minimum for interpretability. \*\* $p < 0.01$

### Corresponding author

Louise Elizabeth Birrell can be contacted at: [l.birrell@unsw.edu.au](mailto:l.birrell@unsw.edu.au)

For instructions on how to order reprints of this article, please visit our website:

[www.emeraldgroupublishing.com/licensing/reprints.htm](http://www.emeraldgroupublishing.com/licensing/reprints.htm)

Or contact us for further details: [permissions@emeraldinsight.com](mailto:permissions@emeraldinsight.com)